

A Cosmic Battery

Ioannis Contopoulos ¹ and Demosthenes Kazanas ²

ABSTRACT

We show that the Poynting–Robertson drag effect in an optically thin advection–dominated accretion flow around active gravitating objects generates strong azimuthal electric currents which give rise to astrophysically significant magnetic fields. Although the mechanism is most effective in accreting compact objects, it seems very promising to also account for the generation of stellar dipolar fields during the late protostellar collapse phase, when the star approaches the main sequence.

Subject headings: MHD — Sun: magnetic fields

1. Introduction

The origin of cosmic magnetic fields remains one of the open questions in astrophysics. Their origin has been sought in the entire history of the Universe going as far back as the QCD phase transition (Sigl, Olinto & Jedamzik 1997) the electroweak phase transition (e.g. Baym, Bodeker & McLerran 1996; Joyce & Shaposhnikov 1997) or even close to the Planck time (e.g. Ratra 1992). Most theories attempt to produce the observed magnetic fields by starting with some ‘seed magnetic field’ and follow its amplification by means of a ‘dynamo mechanism’, i.e. a mechanism which can convert kinetic energy of the conducting fluid, into which the magnetic field is supposedly frozen, into magnetic field energy density (Moffat 1978; Parker 1979). Unfortunately, when one considers the back reaction of the stretched and folded magnetic field on the dynamics of the conducting fluid, dynamo action seems to be very ineffective, in that as the field at the small diffusive scales reaches equipartition, the large-scale component remains several orders of magnitude weaker than astrophysical observed magnetic fields (Vainshtein & Cattaneo 1992).

In addition to these problems with the effectiveness of the dynamo theory, the origin of the ‘seed magnetic field’ needed to be amplified is also an issue, assumed in most cases as a

¹Physics Department, University of Crete, P. O. Box 2208, Heraklion 71003, Greece

²NASA/Goddard Space Flight Center, Greenbelt, MD 20771

boundary condition. However, the consideration that in its initial state the Universe was homogeneous and isotropic to a high degree, excludes the presence of large scale magnetic fields. The seed magnetic field is therefore considered to be generated by some ‘battery mechanism’, the most common being the ‘Biermann battery’ (Biermann 1950), i.e. a mechanism for generating large scale electric currents based on the thermoelectric effect. In this respect it is worth reminding the reader that any battery mechanism depends on the possibility of imparting and sustaining different velocities to the electrons and protons in a plasma, thus creating an electric current. Obviously, in order for such an electric current to give rise to a steady (rather than transient) magnetic field it is necessary that the line integral of the proton-electron velocity difference along a closed, large scale circuit does not vanish. This leads directly to the notion that the generation of magnetic fields should be naturally associated with motions in which the integral $\oint \mathbf{v} \cdot d\mathbf{l}$ does not vanish, i.e. motions with non-zero circulation, a quantity also thought to be zero at the early Universe but generated later, in situations which involve the presence of dissipation, thus suggesting dissipation as a necessary condition for the generation of large scale magnetic fields too.

Herein we revisit the theory of battery generated cosmic magnetic fields by employing a specific mechanism for producing such currents in a specific astrophysical setting, namely the Poynting–Robertson drag effect on the electrons of an Advection–Dominated Accretion Flow (Narayan & Yi 1994, 95; hereafter ADAF) in the vicinity of an accreting object. We examine the time evolution of these magnetic fields in a variety of several astrophysical settings. We find that this effect may be significant enough, at some phase in the evolution of the accreting object, as to not require any further amplification in order to account for the observed, present day values associated with the magnetic fields of the class of objects in question.

In § 2 we give a brief outline of our mechanism and examine qualitatively its implications and effects in different astrophysical settings. In § 3 we integrate numerically the associated equations and compare our results to those derived earlier on qualitative basis. Finally, in § 4 the results are summarized and conclusions are drawn.

2. The model

The model is simple: Consider a point-like astrophysical energy source of luminosity L , and an optically thin, fully ionized ADAF with rotational velocity $v_\phi = Ar\Omega_K$ at distances $r_{\text{in}} \leq r \leq r_{\text{out}}$ around the above energy source (Ω_K is the Keplerian angular velocity, and $A < 1$ is a constant parameter of the flow). Assuming that its scattering depth is sufficiently small that it scatters photons from the center only once, each plasma electron feels an

azimuthal force equal to

$$F_{\text{P-R}} = -\frac{L\sigma_{\text{T}}}{4\pi r^2 c} \frac{v_{\phi}}{c} \quad (1)$$

against its direction of rotation in the region between r_{in} and r_{out} where the plasma is optically thin. This is the Poynting–Robertson radiation drag. The positive ions feel a force $(m_i/m_e)^2$ times smaller which can be neglected. As a consequence of the above force, the electrons tend to lag behind the protons, and an azimuthal electric current density

$$J = \frac{c}{4\pi} \left(\frac{\partial B_r}{\partial z} - \frac{\partial B_z}{\partial r} \right) \sim -\frac{c}{4\pi} \frac{\partial B_z}{\partial r} \quad (2)$$

will develop³⁴. Now, during the ADAF phase, the plasma is turbulent with a turbulent plasma conductivity σ_t (see below), whereas when the ADAF phase ceases, the plasma conductivity attains a much higher value, $\sigma_{\text{Spitzer}} = 10^{17} T_6^{1.5} \text{ s}^{-1}$ (e.g. Zombeck 1992; T_n is the plasma temperature in units of 10^n K). On average, the effect of the finite plasma conductivity σ_{eff} can be parametrized as an extra azimuthal drag force equal to

$$F_{\text{drag}} = \frac{Je}{\sigma_{\text{eff}}} , \quad (3)$$

acting on the electrons (similarly for the ions). The azimuthal component of the microscopic momentum (or force) equation for the electrons can be written as

$$-e(E - \frac{v_r}{c} B_z) + F_{\text{P-R}} + F_{\text{drag}} = 0 , \quad (4)$$

where E is the azimuthal component of the electric field; and v_r is the radial flow velocity; e is taken positive. Combining eqs. (1), (2), (3), and (4) the axial component of the induction equation $\partial B_z / \partial t = -(c/r) \partial(rE) / \partial r$ yields

$$\frac{\partial B_z}{\partial t} = \frac{1}{r} \frac{\partial}{\partial r} \left\{ \frac{L\sigma_{\text{T}} A \Omega_{\text{K}}}{4\pi c e} - r v_r B_z + \frac{c^2}{4\pi \sigma_{\text{eff}}} r \frac{\partial B_z}{\partial r} \right\} , \quad (5)$$

³In order to keep the analysis simple, we will assume a cylindrical distribution for the magnetic field inside the geometrically thick ADAF, i.e. we will neglect its radial component and the associated magnetic field tension.

⁴The reader should not worry that ‘flux-freezing’ is not satisfied as electrons and ions move at slightly different azimuthal velocities. Electric currents generate magnetic forces which balance the forces $\mathbf{F}_e$ and $\mathbf{F}_i$ acting on electrons and ions respectively. The reader can convince him/herself that when the problem is solved, the two plasma constituents perform $\mathbf{F}_{e,i} \times \mathbf{B} / (q_{e,i} B^2)$ drifts across the above force fields, i.e. they do not move together even under conditions of flux-freezing (this analysis remains outside the main scope of the present work, and can be found in Ciolek & Mouschovias 1993).

in the range of radii $(r_{\text{in}}, r_{\text{out}})$. Ψ is defined through the relation $B_z = r^{-1} \partial \Psi / \partial r$, and $2\pi \dot{\Psi}$ is the growth rate of the total magnetic flux contained within radius r . One should note that eq. (5) includes both the advection of magnetic field through the term $-rv_r B_z$, as well as its diffusion through the term $c^2 / 4\pi \sigma_{\text{eff}} (r \partial B / \partial r)$. The connection between magnetic field and circulation discussed in the introduction, then, suggests that in an accretion disk, the turbulent magnetic diffusivity should also be proportional to the quantity effecting the dissipation in the accretion disk, namely the turbulent viscosity parameter. Thus one may write (e.g. Reyes–Ruiz & Stepinski 1996)

$$\eta_{\text{eff}} \equiv c^2 / (4\pi \sigma_{\text{eff}}) \sim \mathcal{P}_m \nu \sim \mathcal{P}_m \alpha \xi r^2 \Omega_K, \quad (6)$$

where, $\alpha < 1$ (the usual Shakura–Sunyaev (1973) α parameter) and $\xi < 1$ (the ratio of the local flow temperature to the virial temperature) are taken to be constant flow parameters (we remind the reader that $A \equiv \Omega / \Omega_K = (1 - 5\xi/2 - \alpha^2 \xi^2/2)^{1/2}$; Narayan & Yi 1994, 95); $\mathcal{P}_m$ is the ratio of magnetic diffusivity to viscosity, the so-called magnetic Prandtl number. Note in addition that in an ADAF, the radial velocity can be related to the azimuthal one through the relation

$$v_r \sim -\alpha \xi r \Omega_K \sim -\eta_{\text{eff}} / (\mathcal{P}_m r). \quad (7)$$

Substituting all the above in the induction equation leads to

$$\frac{\partial B_z}{\partial t} = \frac{1}{r} \frac{\partial}{\partial r} \left\{ \frac{L \sigma_T A \Omega_K}{4\pi c e} + r^2 \Omega_K \alpha \xi \left[B_z + \mathcal{P}_m r \frac{\partial B_z}{\partial r} \right] \right\} \equiv \frac{1}{r} \frac{\partial \dot{\Psi}}{\partial r} \quad (8)$$

The r.h.s. of eq. (8) consists of a source term (independent of σ_{eff}), an advection term, and a diffusion term. The source term is the Poynting–Robertson azimuthal electric current and leads to field growth which is fastest at $r = r_{\text{in}}$. This gives rise to a distribution of closed poloidal magnetic field loops, with their center somewhere inside r_{in} . As we will see in the next subsection, *when $\mathcal{P}_m \gtrsim 2$, the magnetic flux contained inside r_{in} grows almost linearly in time, as long as the ADAF persists. When rotation and accretion cease, and $\sigma_{\text{eff}} \sim \sigma_{\text{Spitzer}}$, this flux will remain effectively trapped in the plasma over timescales*

$$\tau \simeq \frac{r_{\text{in}}^2}{\eta_{\text{Spitzer}}} \simeq 4x^2 M_0^2 T_6^{1.5} \text{ yr}. \quad (9)$$

M_n is the mass of the central object in units of $10^n M_\odot$; $x \equiv r/r_S$ is the radial coordinate in units of the central object’s Schwarzschild radius r_S .

Obviously, this mechanism will cease being effective when the magnetic field becomes strong enough to affect the dynamics of accretion. In addition, a magnetic field which becomes dynamically important will also affect the torques from the ADAF onto the star

(or vice versa). We prefer to defer discussion of the implications of this aspect of the problem and the rotational evolution of the compact object to a future publication; as indicated by the present analysis these could indeed be significant.

It is of interest to examine the time scales necessary to accumulate enough flux via the above mechanism in the most optimistic case with $\mathcal{P}_m > 2$, for the corresponding magnetic field to affect significantly the dynamics of accretion, as this may be an important component to the formation of jets, which are ubiquitous in accreting compact objects. Assuming an almost constant growth rate of magnetic flux Ψ of the order of the first term in the r.h.s. of eq. (5),

$$\Psi \sim 3 \cdot 10^{23} \frac{f}{x^{3/2}} \left(\frac{t}{\text{yrs}} \right) \text{ G cm}^2, \quad (10)$$

where, f is the fraction of the source luminosity compared to the Eddington luminosity. One should note that the latter expression is independent of the mass of the system and depends only on its luminosity through the parameter f (for compact objects x is considered to be of order unity). However, the magnetic field itself is bounded by considerations associated with equipartition of the magnetic energy density with that of the ram pressure of almost free-falling matter,

$$B_{eq} \sim 9 \cdot 10^7 x^{-1.25} \dot{m}^{0.5} M_0^{-0.5} \text{ G}, \quad (11)$$

where $\dot{m}$ is the accretion rate in units of the Eddington rate. Following Narayan & Yi 1994, we took $\alpha = 0.3$, $\xi = 0.38$, and $A = 0.2$. Assuming further that $B(r \leq r_{\text{in}}) \sim \Psi(r_{\text{in}})/r_{\text{in}}^2$, and that $f \simeq \dot{m}/x$, one obtains

$$\tau_{eq} \sim 4 \cdot 10^{-5} x^{3.25} M_0^{1.5} \dot{m}^{-0.5} \text{ yr}. \quad (12)$$

This value is comparable to the Eddington time ($\sim 10^8$ yrs) for an AGN with a massive central black hole ($M_9 \sim 1$), but decreases very quickly with the decrease of the mass.

3. Numerical Simulation

The detailed time evolution of the magnetic field can be computed by direct integration of the time evolution equation (5). We introduce the dimensionless variables

$$\hat{r} = r/r_{\text{in}}, \quad \hat{t} = t/t_o, \quad b = B/B_o, \quad \psi = \dot{\Psi}t_o/(r_{\text{in}}^2 B_o), \quad (13)$$

where, $t_o \equiv (\Omega_K(r_{\text{in}})\alpha\xi)^{-1}$; $B_o \equiv L\sigma_T A/(4\pi c e r_{\text{in}}^2 \alpha\xi)$. The induction equation can now be rewritten in dimensionless form applicable in various astrophysical situations,

$$\frac{\partial b}{\partial \hat{t}} = \frac{1}{\hat{r}} \frac{\partial}{\partial \hat{r}} \left\{ \hat{r}^{-1.5} + \hat{r}^{0.5} \left[b + \mathcal{P}_m \hat{r} \frac{\partial b}{\partial \hat{r}} \right] \right\} \equiv \frac{1}{\hat{r}} \frac{\partial \psi}{\partial \hat{r}}. \quad (14)$$

The induction equation is integrated numerically in the interval $1 \leq \hat{r} \leq \hat{r}_{\text{out}}$. In order to keep the number of free parameters to a minimum, we take $\hat{r}_{\text{out}} \gg 1$. We follow the numerical procedure described in the Appendix of Mouschovias & Morton (1991), and implement the ‘free boundary’ condition $\partial b / \partial \hat{r} = 0$ at $\hat{r} = \hat{r}_{\text{in}}$, allowing thus any amount of magnetic flux that would be advected inside $\hat{r}_{\text{in}}$ to enter unimpeded. We discovered that, for magnetic Prandtl numbers $\mathcal{P}_m > 2$, the dimensionless magnetic flux inside dimensionless radius $\hat{r} = 1$ rises almost linearly with time, with a dimensionless flux growth rate $\dot{\psi}$ of order unity. In fact, when $\mathcal{P}_m \geq 2$, eq. (14) has the steady-state solution

$$b(\hat{r}) = \frac{1}{(2\mathcal{P}_m - 1)} \left[\frac{1}{\hat{r}^2} - \frac{4}{\hat{r}^{0.5}} \right], \quad (15)$$

with, $\dot{\psi} = 2(\mathcal{P}_m - 2)/(2\mathcal{P}_m - 1)$. We understand this interesting behavior as follows: The Poynting–Robertson effect generates an azimuthal electric current resulting in closed loops of poloidal magnetic field with a distribution of positive magnetic flux inside r_{in} (where the magnetic field points, say, upward) and negative flux outside, i.e. in the regions with opposite field polarity. The rate of positive flux growth inside r_{in} is given by the first term in the r.h.s. of eq. (14). When $\mathcal{P}_m \geq 2$, diffusion is dominant, and the distribution of *negative* flux outside r_{in} diffuses outward to larger distances. As a result, positive flux is generated without limit (except of course for the equipartition limit) in the interior. On the other hand, when $\mathcal{P}_m < 2$, the inward advection of the negative flux dominates over its outward diffusion. As a result, the steady-state positive flux accumulated inside r_{in} remains finite, since its rate of growth by the Poynting–Robertson mechanism is balanced by the rate of inward advection of the return (negative) flux.

In figure 1 we exhibit the spatio-temporal evolution of the magnetic field and the associated flux resulting from our numerical solution to eq. (14). In figure 1a we show the evolution of the radial distribution of the magnetic field as a function of the dimensionless radius r/r_{in} for $\mathcal{P}_m = 2.5$. Each curve corresponds to an increment in time by $10t_o$. In figure 1b we exhibit the evolution of the magnetic flux held within r_{in} as a function of time for different values of $\mathcal{P}_m$. The reader should note that according to eq. (15), for Prandtl numbers $\mathcal{P}_m > 2$, a steady state situation, though it corresponds to a constant value for the magnetic field in the region $r/r_{\text{in}} > 1$, demands a linear increase for the magnetic flux Ψ for $r/r_{\text{in}} < 1$, with the return flux diffusing steadily to infinity.

In table 1, we introduce dimensions, and apply our mechanism to specific astrophysical situations. The mechanism is most effective at short scales, especially around compact objects (neutron stars, black holes of both stellar and galactic sizes). One could also suggest that magnetic flux generated quickly around active galactic nuclei, could be carried by winds out to galactic (and even extragalactic) scales (similar mechanisms have already been

proposed in order to account for the seed galactic dynamo field, e.g. Chakrabarti, Rosner, Vainshtein 1994).

When we apply our mechanism to the most interesting stellar case, namely that of the Sun, we consider a scenario in which the innermost, optically thick, hydrogen burning core grows toward its main sequence values $r_c \sim 0.1 r_\odot$, and $M_c \sim 0.1 M_\odot$ respectively, or $x \simeq 2 \times 10^5$. We then consider that the remainder of the stellar mass ($\simeq 1 M_\odot$) accretes onto this component in the form of an ADAF, releasing and radiating away its gravitational energy. In this respect, the recent discovery of X-ray emission from objects in this class (Feigelson, private communication) is, to say the least, encouraging. Finally, we assume that the newly forming core radiates at some fraction $f = 0.1$ of the Eddington luminosity that corresponds to a mass equal to $0.1 M_\odot$ (or equivalently $L = 332 L_\odot$; Casanova *et al.* 1995; Gross *et al.* 1998). The above are supposed to be crude approximations during the final stages of almost free-fall protostellar collapse. In the general protostellar collapse phase,

$$\Psi \simeq 0.4 \Psi_\odot f M_{-1}^{1.5} (r_c / 0.1 r_\odot)^{-1.5} (t / 10^6 \text{ yr}) . \quad (16)$$

Note that because of the linearity between the mass and the radius of the initial (proto)stellar cores, the above estimate, in accordance with eq.(10), is independent of their precise values, depending only on their ratio x . As one can see, for $f \sim 0.1$, it takes several million years to build the present day total observed solar dipolar magnetic flux $\Psi_\odot = 1 \text{ G} \cdot r_\odot^2$. Considering the simplicity of these arguments, the agreement with observation is not unreasonable.

We would like to discuss what is the geometry of the field thus generated and its fate after the mechanism ceases to be important (i.e. after the optically thin accretion flow dies out): The field is contained at the very core, in a layer around r_c . In the absence of turbulence, the diffusion time-scale is of the order of several billion years, and the field will

	r_{in}	L	M	τ_{eq}	B_{eq}
Black Hole	$3 \cdot 10^6$	10^4	10	$7 \cdot 10^{-4}$	$5 \cdot 10^7$
Neutron Star	10^6	10^4	1	10^{-3}	$2 \cdot 10^7$
White Dwarf	10^8	10	1	$2 \cdot 10^4$	$2 \cdot 10^4$
Protosun	$7 \cdot 10^9$	332	10^{-1}	$2 \cdot 10^9$	$8 \cdot 10^3$
AGN	$3 \cdot 10^{13}$	10^{12}	10^8	$7 \cdot 10^7$	$5 \cdot 10^3$
Galaxy	10^{22}	10^{10}	10^{10}	$6 \cdot 10^{30}$	$6 \cdot 10^{-5}$

Table 1: r_{in} in cm; L in $L_\odot$; M in $M_\odot$; τ_{eq} in yr; $B_{eq} \equiv \Psi_{eq}/r_{\text{in}}^2$ in G.

remain effectively trapped inside the solar radiation zone, i.e. below the turbulent surface convection zone. Any small fraction of this field which diffuses out of the sun will assume an overall dipolar geometry at large scales. In our mechanism the field is generated in situ, rather than being brought in from large distances. This magnetic field geometry should naturally interact with stellar differential rotation, leading thus to torsional oscillations which might account for the solar cycle (e.g. Layzer, Rosner & Doyle 1979; Goode & Dziembowski 1991).

4. Discussion

The scenario described in this paper is not new. Previous researchers considered the rotation of solar coronal layers (Cattani & Sacchi 1966) as the source of the solar magnetic field, and the rotation of protogalaxies with respect to the cosmic microwave background radiation (Harrison 1970; Mishustin & Ruzmaikin 1972), as a means for producing the observed galactic field and (correctly) obtained very weak magnetic field values in both cases. However, to the best of our knowledge, this is the first time that this scenario has been applied to other classes of objects, in particular to optically thin accretion flows onto compact objects, where it appears to have its greatest impact, as the figures in table 1 indicate.

Concerning the galactic magnetic fields, or better the value of the galactic magnetic flux $\Psi_g \sim 3 \times 10^{36} \text{ G cm}^2$ (this is mainly azimuthal and confined on the plane of the galaxy) which presents the largest discord between the predictions of our model and observations (see table 1), one could conceivably consider its generation during an early AGN phase of our galaxy, instead of the era of galaxy formation; as such one could argue for the generation of the requisite flux near a massive black hole at the galactic center and its diffusion to large distances as discussed in the previous section, or as invoked by Chakrabarti, Rosner & Vainshtein (1994). Assuming that such an AGN stage lasted $\sim 10^9$ yrs, i.e. a few times the Eddington time (a quantity which is independent of the mass of the compact object), during which the mass galactic center black hole presumably grew to its present size and the associated magnetic field reached its equipartition value, eq.(10) yields a poloidal flux roughly 10^4 times smaller than that observed. However, the galactic differential rotation in the ensuing 10^{10} years (corresponding to roughly 100 galactic rotations at the outer regions of the galaxy and 100 - 300 times this value for its inner regions) could easily amplify this figure to its presently observed value!

Considering in particular the generation of the fields in stars, this scenario can yield observationally interesting values for the magnetic field only if one is willing to consider

that during the pre-main sequence collapse of the cloud out of which the star formed, an advection-dominated accretion flow was probably present, and only if one is willing to accept values for the luminosity of this object (as given by f) and the duration of this process similar to those given above. However, even for lower values for f or for times shorter than $\sim 10^7$ years, as in the above galactic case, the magnetic flux in the stellar interior could subsequently be amplified linearly with time by the interior differential rotation to observationally interesting values. Given the very general scope of this work we would not like to pursue these issues beyond this level at present. We feel however that the assumptions about the pre-main sequence protostellar evolution employed in our arguments are not unreasonable. Interestingly enough, were the stellar magnetic field to be generated as discussed in the previous section, it should be strong and largely confined to the deep stellar interior, leading naturally to the torsional oscillations between differentially rotating stellar layers which have been purported to account for stellar cycles.

One cannot fail to notice that, when the outside magnetosphere can support electric currents (i.e. when there is no vacuum outside), the magnetic field geometry obtained in our numerical simulations, leads naturally to the electromagnetic interaction between the central region and the extended disk. The reason is that the differential rotation ‘twists’ the newly formed field, generating thus magnetic torques on the central part. The magnetic field twisting leads to another very important effect, namely the overall opening of the closed loop geometry (Newman, Newman & Lovelace 1992; Goodson, Winglee & Böhm 1997).⁵ As one can see in the above references, the axial part of the expanding plasma + magnetic field remains well collimated. Under certain physical circumstances (Shibata, Tajima & Matsumoto 1992; Matsumoto, Matsuzaki, Tajima & Shibata 1997), this effect might also proceed explosively, leading thus to the expulsion of a significant fraction of the disk material in the form of collimated fast axially moving plasma outflows (the astrophysical plasma gun, Contopoulos 1995; Matsumoto *et al.* 1996).

In this respect it is of interest that the time necessary for the magnetic field to become dynamically important is larger than the Eddington time for the most massive AGN with $M \gtrsim 3 \cdot 10^8 M_\odot$. Thus one could speculate that the bright, optically selected QSOs have not had yet time to generate dynamically significant amounts of magnetic flux. One could further speculate that when they do so, the associated magnetic field would become an important component of the dynamics of accretion, facilitating the production of radio jets and leading to conversion of these objects from radio quiet to radio loud AGN.

⁵In a simple ‘mechanistic’ interpretation, this can be thought of as the dynamical spring-like release through unwinding, of the azimuthal magnetic field generated by the differential rotation of the magnetic field footpoints.

The evolution of AGN and in particular its possible relation to their radio properties are far from being understood. The arguments put forward herein, associated with the generation and evolution of their magnetic fields may offer a new way of looking into this problem. Interestingly, the models of Falcke & Biermann (1995) which provide scalings of the radio loud objects from the quasar to the galactic binary source domain, do depend, to a large extent, on equipartition arguments such as those discussed above. It would be worthwhile to examine whether these arguments could be put in the context of an evolutionary scenario of both the mass or luminosity and the associated magnetic field.

To conclude, it seems that the mechanism discussed above enters as a new, previously unaccounted for, player in the ongoing studies of pre-main sequence stellar collapse, the generation of stellar and galactic magnetic fields, the spin evolution of accreting magnetic stars, and the generation of magnetically driven jets.

We would like to thank the anonymous referee for his constructive, penetrating criticism which, inducing a major revision to an earlier version of this work, contributed significantly to its improvement. D.K. would like to acknowledge an informative discussion with E. Feigelson. I.C. wishes to acknowledge support by grant 107526 of the General Secretariat of Research and Technology of Greece.

REFERENCES

- Baym, G., Bodeker, D. & McLerran, L., 1996, Phys. Rev. D53, 662
- Biermann, L., 1950, Z. Naturforsch, 5a, 65
- Casanova, S., Montmerle, T., Feigelson, E. D. & André, P., 1995, ApJ, 439, 752
- Cattani, D. & Sacchi, C., 1966, Nuovo Cimento, 46B, 8046
- Chakrabarti, S. K., Rosner, R. & Vainshtein, S. I., 1994, Nature, 368, 434
- Ciolek, G. E. & Mouschovias, T. Ch., 1993, ApJ, 418, 774
- Contopoulos, J., 1995, ApJ, 450, 616
- Falcke, H. & Biermann, P. L., 1995, A& A, 293, 665
- Goode, P. R. & Dziembowski, W. A., 1991, Nature, 349, 223
- Goodson, A. P., Winglee, R. M. & Böhm, K-H, 1997, ApJ, 489, 199
- Grosso, N., Montmerle, T., André, P., Casanova, S., Feigelson, E. D. & Gregorio-Hetem, J., 1998, in ‘Star Formation with the ISO’, ASP Conf. Series, 132, 354

- Harrison, E. R., 1970, MNRAS, 147, 279
- Joyce, M. & Shaposhnikov, M., 1997, Phys. Rev. Lett., 79, 1193
- Layzer, D., Rosner, R. & Doyle, H. T., 1979, ApJ, 229, 1126
- Matsumoto, R., Matsuzaki, T., Tajima, T. & Shibata, K., 1997, BAAS, 191, 27.11
- Matsumoto, R., Uchida, Y., Hirose, S., Shibata, K., Hayashi, M. R., Ferrari, A., Bodo, G. & Norman, C., 1996, ApJ, 461, 115
- Mishustin, I. N. & Ruzmaikin, A. A., 1972, Sov. Phys. JETP, 34, 233
- Moffatt, H. K., 1978, Magnetic Field Generation in Electrically Conducting Fluids (Cambridge: Cambridge Univ. Press)
- Mouschovias, T. Ch. & Morton, S. A., 1991, ApJ, 371, 296
- Narayan, R. & Yi, I., 1994, ApJ, 428, L13
- Narayan, R. & Yi, I., 1995, ApJ, 452, 710
- Newman, W. I., Newman, A. L. & Lovelace, R. V. E., 1992, ApJ, 392, 699
- Parker, E. N., 1979, Cosmical Magnetic Fields (Oxford: Clarendon)
- Ratra, V., 1992, ApJ, 391, L1
- Reyes–Ruiz, M. & Stepinski, T. F., 1996, ApJ, 459, 653
- Shakura, N. I. & Sunyaev, R. A., 1973, A& A, 24, 337
- Shibata, K., Tajima, T. & Matsumoto, R., 1990, ApJ, 350, 295
- Sigl, G., Olinto, A. V. & Jedamzik, 1997, Phys. Rev D55, 5582
- Vainshtein, S. I. & Cattaneo, F., 1992, ApJ, 393, 165
- Zombeck, M. V., 1992, Handbook of Space Astronomy & Astrophysics, 2nd ed. (Cambridge: Cambridge Univ. Press)

Figure Captions

Fig. 1.— Magnetic field evolution across the rotating plasma. In (a), we plot the magnetic field $b = B_z/B_o$ distribution as a function of the dimensionless radius r/r_{in} , at time intervals $10t_o$, for $\mathcal{P}_m = 2.5$. In (b), we plot the growth of the magnetic flux $\psi = \Psi/(r_{\text{in}}^2 B_o)$ contained inside $r = r_{\text{in}}$ with time, for different Prandtl numbers $\mathcal{P}_m$. One sees clearly the fast growth of the field around r_{in} , and the almost linear growth of the central magnetic flux for $\mathcal{P}_m > 2$. The present (kinematic) integration should not be continued beyond τ_{eq} , when the field becomes dynamically important.





Fig. 1.—